

Non-Fatal Self-Harm in Ireland - Findings from the National Registry of Deliberate Self-Harm, Annual Report 2012

Griffin, E.,¹
Arensman, E.,²
Wall, A.,³
Corcoran, P.,⁴
Perry, I.J.⁵
National Suicide
Research
Foundation,
University College
Cork^{1,2,3}
Department of
Epidemiology and
Public Health,
University
College
Cork^{2,4,5}

ABSTRACT

The eleventh annual report from the National Registry of Deliberate Self-Harm is based on data collected on persons presenting to all 35 hospital emergency departments as a result of self-harm in 2012 in the Republic of Ireland. In 2012, the Registry recorded 12,010 presentations to hospital due to deliberate self-harm nationally, involving 9,483 individuals. Taking the population into account, the age-standardised rate of individuals presenting to hospital following deliberate self-harm in 2012 was 211 per 100,000, a 2% decrease on the rate in 2011. However, the rate in 2012 was still 12% higher than that in 2007. In 2012, the national male rate of deliberate self-harm was 195 per 100,000, and the female rate of deliberate self-harm in 2012 was 228 per 100,000. Despite the overall decrease in 2012, the male rate has increased by 20% since 2007 and the female rate has increased by 6% over the same period (Table 1).

Table 1 - Number of Deliberate Self-Harm Presentations and Persons who Presented in the Republic of Ireland in 2002-2012

Year	Presentations		Persons	
	Number	%Diff	Number	%Diff
2002	10,537	-	8,421	-
2003	11,204	+6%	8,805	+5%
2004	11,092	-1%	8,610	-2%
2005	10,789	-3%	8,594	<-1%
2006	10,688	-1%	8,218	-4%
2007	11,084	+4%	8,598	+5%
2008	11,700	+6%	9,218	+7%
2009	11,966	+2%	9,493	+3%
2010	12,337	+3%	9,887	+4%
2011	12,216	-1%	9,834	<-1%
2012	12,010	-2%	9483	-4%

Despite a decrease in the number of self-harm presentations in 2012 from 2011, the proportion accounted for by repetition in 2012 (21.0%) was higher than that in 2010 or 2011, and similar to the years 2003-2009 (range: 20.5-23.1%). This confirms that repetition continues to pose a major challenge to hospital staff and family members involved.

Drug overdose was the most common method of self-harm, involved in 69% of all acts registered in 2012, and more so in women (75%) than in men (62%). Minor tranquilisers, paracetamol-containing medicines and anti-depressants/mood stabilisers were involved in 41%, 28% and 22% of drug overdose acts. Attempted hanging was

involved in 7% of all deliberate self-harm presentations (10% for men and 3% for women). At 776, the number of presentations involving attempted hanging has increased significantly by 6% from 2011 and by 75% from 2007 (n=444). This is the greatest number of deliberate self-harm presentations involving hanging recorded by the Registry and is 75% higher than the number recorded in 2007 (n=444). Cutting was the only other common method of self-harm, involved in 23% of all episodes and was significantly more common in men (26%) than women (21%). Alcohol was involved in 38% of all cases. While overall alcohol involvement decreased slightly from 2011, alcohol was significantly more often involved in male episodes of self-harm than female episodes (42% versus 36%, respectively).

Overall, in 12% of 2012 cases, the patient left the emergency department before a next care recommendation could be made. Following their treatment in the emergency department, inpatient admission was the next stage of care recommended for 38% of cases, irrespective of whether general or psychiatric admission was intended and whether the patient refused or not. Of all deliberate self-harm cases, 28% resulted in admission to a ward of the treating hospital whereas 10% were admitted for psychiatric inpatient treatment from the emergency department.

Following successive increases in deliberate self-harm in Ireland during the period 2007-2010, the 2012 Annual Report of the National Registry of Deliberate Self-Harm shows a second subsequent annual decrease. Considering the relatively small reduction, this should be interpreted with caution since it would be premature to conclude that this indicates a decreasing trend. The 2012 Registry outcomes underline an ongoing need for prevention and intervention programmes to be implemented at national level. Increased and continued support should be provided for evidence-based and best practice prevention and mental health promotion programmes in line with priorities in Reach Out, National Strategy for Action on Suicide Prevention (2005-2014) and Vision for Change, the Report of the Expert Group on Mental Health Policy.

SOURCE

Griffin, E. Arensman, E. Wall, A. Corcoran, P. Perry, I.J. (2013). National Registry of Deliberate Self-Harm Annual Report 2012. Cork: National Suicide Research Foundation.

FUNDING

The National Registry of Deliberate Self-Harm is funded by the Health Service Executive's National Office for Suicide Prevention.